University of California Agriculture and Natural Resources

Kern UCCE/DWR Weekly Crop Water Use Report

Making a Difference for California

WEEKLY SOIL MOISTURE LOSS IN INCHES

(Estimated Crop Evapotranspiration or ETc) 03/14/16 through 03/20/16

Crops (Leafout Date)	#5 Shafter			#125 Arvin-Edison			#146 Belridge			
	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's	Past Week	Accum'd	Next Week's	
	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated	of Water	Seasonal	Estimated	ì
	Use	Water Use	ETc	Use	Water Use	ETc	Use	Water Use	ETc	
Almonds (2/22) *	0.73	2.19	0.75	0.76	2.37	0.82	0.79	2.35	0.86	i
Pistachio * **	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	i
Citrus	0.69	4.37	0.68	0.70	4.71	0.76	0.75	4.84	0.77	ı
Grapes (3/10) (late season table, 75% cover)	0.13	0.22	0.16	0.13	0.24	0.17	0.14	0.25	0.17	1
Winegrapes (3/10) (50% cover) ***	0.13	0.22	0.14	0.13	0.22	0.14	0.13	0.23	0.18	1
Alfalfa	0.98	6.03	0.96	1.01	6.40	1.04	1.06	6.64	1.09	i
Cotton	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	i.
Past 7 days precipitation (inches)		0.10			0.00			0.00		
Accumulated precipitation (inches)		2.18			4.40			2.54		

Accumulations started on February 22, 2016 or on the approximate leafout date for a specific orchard crop as indicated in parentheses. Criteria for beginning this report are based on the season's last significant rainfall event where the soil moisture profile is estimated to be near its highest level for the new season.

^{***} Winegrapes irrigated at 50% of ETo starting June 1 to end of September.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹													
Crops		#5 Shafter				#125 Arvin-Edison				#146 Belridge			
System Efficiency >>	65%	75%	85%	95%	65%	75%	85%	95%	65%	75%	85%	95%	
Almonds (2/22)	1.2	1.0	0.9	0.8	1.2	1.0	0.9	0.8	1.2	1.1	0.9	0.8	
Pistachio	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
Citrus	1.1	0.9	0.8	0.7	1.1	0.9	0.8	0.7	1.1	1.0	0.9	0.8	
Grapes (3/10) (late season table, 75% cover)	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	
Winegrapes (3/10) (50% cover)	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	0.2	0.2	0.2	0.1	
Alfalfa	1.6	1.3	1.2	1.0	1.6	1.3	1.2	1.1	1.6	1.4	1.2	1.1	
Cotton	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%

For further information concerning all counties receiving this report, contact the Kern Co. Farm Advisor's office at (661) 868-6218.

^{*} Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

^{**} Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 - resulting in about 8% greater water use than shown in these tables.