

WEEKLY CROP WATER USE - Based on local CIMIS Weather Stations (in inches)
(Estimated Crop Evapotranspiration or ETc)
09/24/18 through 09/30/18

Crops (Leafout Date)	#5 Shafter				#125 Arvin-Edison				#146 Belridge		
	9/17 - 9/23 Water Use	Accum'd Seasonal Water Use	9/24 - 9/30 Estimated ETc		9/17 - 9/23 Water Use	Accum'd Seasonal Water Use	9/24 - 9/30 Estimated ETc		9/17 - 9/23 Water Use	Accum'd Seasonal Water Use	9/24 - 9/30 Estimated ETc
Almonds (3/7) *	1.44	44.06	1.18		1.58	48.17	1.27		1.34	42.61	1.19
Pistachio (4/15) * **	1.51	34.71	1.28		1.65	38.64	1.37		1.41	33.07	1.29
Citrus (2/1)	1.06	33.70	0.75		1.14	36.29	0.82		0.97	33.10	0.76
Grapes (3/20) (late season table, 75% cover)	1.93	42.67	1.62		2.10	47.46	1.75		1.77	40.61	1.63
Winegrapes (3/20)(50% cover) ***Irrigation Finished	0.70	22.01	0.60		0.78	23.98	0.64		0.64	21.35	0.60
Alfalfa (2/1)	1.37	47.94	1.16		1.51	52.01	1.25		1.27	46.99	1.17
Cotton (4/15) Irrigation Finished	1.27	32.22	0.93		1.41	36.29	1.02		1.18	30.45	0.94
Past 7 days precipitation (inches)	0.00				0.00				0.00		
Accumulated precipitation (inches) (1/1/18)	3.66				4.76				3.31		

Accumulations started on the approximate leafout date for a specific orchard crop as indicated in parentheses. Criteria for beginning this report are based on the season's last significant rainfall event where the soil moisture profile is estimated to be near its highest level for the new season.

* Estimates are for orchard floor conditions where vegetation is managed by some combination of strip applications of herbicides, frequent mowing or tillage, and by mid and late season shading and water stress. Weekly estimates of soil moisture loss can be as much as 25 percent higher in orchards where cover crops are planted and managed more intensively for maximum growth.

** Very vigorous, non-salt affected peak season pistachio Kc can be as high as 1.19 – resulting in about 8% greater water use than shown in these tables.

*** Winegrapes irrigated at 50% of ETo starting June 1 to end of September.

PAST WEEKLY APPLIED WATER IN INCHES, ADJUSTED FOR EFFICIENCY ¹

Crops	#5 Shafter					#125 Arvin-Edison					#146 Belridge			
	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
System Efficiency >>	65%	75%	85%	95%		65%	75%	85%	95%		65%	75%	85%	95%
Almonds (3/7)	2.2	1.9	1.7	1.5		2.4	2.1	1.9	1.7		2.1	1.8	1.6	1.4
Pistachio (4/15)	2.3	2.0	1.8	1.6		2.5	2.2	1.9	1.7		2.2	1.9	1.7	1.5
Citrus (2/1)	1.6	1.4	1.2	1.1		1.8	1.5	1.3	1.2		1.5	1.3	1.1	1.0
Grapes (3/20) (late season table, 75% cover)	3.0	2.6	2.3	2.0		3.2	2.8	2.5	2.2		2.7	2.4	2.1	1.9
Winegrapes (3/20) (50% cover)	0	0	0	0		0	0	0	0		0	0	0	0
Alfalfa (2/1)	2.1	1.8	1.6	1.4		2.3	2.0	1.8	1.6		2.0	1.7	1.5	1.3
Cotton (4/15)	0	0	0	0		0	0	0	0		0	0	0	0

¹ The amount of water required by a specific irrigation system to satisfy evapotranspiration. Typical ranges in irrigation system efficiency are: Drip, 80%-95%; Micro-sprinkler, 80%-90%; Sprinkler, 70%-85%; and Border-furrow, 50%-75%.

For further information concerning all counties receiving this report, contact the Kern Co. Farm Advisor's office at (661) 868-6218.